Assignment 7

1. Find the Fourier transform of $PV\frac{1}{x}$. You may use the results of previous exercises.

2. Show that $\phi_n \to \phi$ implies $\widehat{\phi}_n \to \widehat{\phi}$, where convergence is in the sense of the Schwartz space.

- **3.** Give a full proof of Proposition 6.4: For all multi-indices α ,
- (a) $\partial^{\alpha} \widehat{T} = (-2\pi i)^{\alpha} \widehat{x^{\alpha} T}.$

(b)
$$\hat{\partial}^{\alpha} \tilde{T} = (2\pi \mathrm{i} \, \mathrm{k})^{\alpha} \, \widehat{\mathrm{T}}.$$

- **4.** Consider the function $f(x) = \log |x|$ on \mathbb{R} .
 - (a) Show that f is locally integrable. In case you need the primitive of $\log x$, try to differentiate $x \log x$, that may help!
 - (b) Show that the distributional derivative of $\log |x|$ is $PV_{\frac{1}{x}}^{\frac{1}{x}}$.
- 5. Consider the *Airy equation* in one dimension

$$u'' - xu = 0.$$

A solution is the Airy function $\operatorname{Ai}(x) = \frac{1}{\pi} \int_0^\infty \cos(\frac{1}{3}t^3 + xt) dt$. It is named after the British George Biddell Airy who became Astronomer Royal in 1835.

- (a) Check formally that $\operatorname{Ai}(x)$ is solution to the Airy equation. Hint: differentiate under the integral sign; the primitive of $(t^2 + x) \cos(\frac{1}{3}t^3 + xt)$ is easily found.
- (b) Let

$$u(x) = c \int_{-\infty}^{\infty} e^{ik^3/3 + ikx} dk$$

Give u a rigorous meaning as the Fourier transform of a distribution. Prove that the distribution u satisfies the Airy equation. Hint: take the Fourier transform of the equation.